



AP[®] Physics B

2002 Sample Student Responses


Form B

The materials included in these files are intended for use by AP teachers for course and exam preparation in the classroom; permission for any other use must be sought from the Advanced Placement Program[®]. Teachers may reproduce them, in whole or in part, in limited quantities, for face-to-face teaching purposes but may not mass distribute the materials, electronically or otherwise. These materials and any copies made of them may not be resold, and the copyright notices must be retained as they appear here. This permission does not apply to any third-party copyrights contained herein.

These materials were produced by Educational Testing Service[®] (ETS[®]), which develops and administers the examinations of the Advanced Placement Program for the College Board. The College Board and Educational Testing Service (ETS) are dedicated to the principle of equal opportunity, and their programs, services, and employment policies are guided by that principle.

The College Board is a national nonprofit membership association dedicated to preparing, inspiring, and connecting students to college and opportunity. Founded in 1900, the association is composed of more than 4,200 schools, colleges, universities, and other educational organizations. Each year, the College Board serves over three million students and their parents, 22,000 high schools, and 3,500 colleges, through major programs and services in college admission, guidance, assessment, financial aid, enrollment, and teaching and learning. Among its best-known programs are the SAT[®], the PSAT/NMSQT[®], and the Advanced Placement Program[®] (AP[®]). The College Board is committed to the principles of equity and excellence, and that commitment is embodied in all of its programs, services, activities, and concerns.


Copyright © 2002 by College Entrance Examination Board. All rights reserved. College Board, Advanced Placement Program, AP, SAT, and the acorn logo are registered trademarks of the College Entrance Examination Board. APIEL is a trademark owned by the College Entrance Examination Board. PSAT/NMSQT is a registered trademark jointly owned by the College Entrance Examination Board and the National Merit Scholarship Corporation. Educational Testing Service and ETS are registered trademarks of Educational Testing Service.



2. (15 points)

A ball attached to a string of length ℓ swings in a horizontal circle, as shown above, with a constant speed. The string makes an angle θ with the vertical, and T is the magnitude of the tension in the string. Express your answers to the following in terms of the given quantities and fundamental constants.

- (a) On the figure below, draw and label vectors to represent all the forces acting on the ball when it is at the position shown in the diagram. The lengths of the vectors should be consistent with the relative magnitudes of the forces.



- (b) Determine the mass of the ball.

TENSION = $= T \cos \theta$

SINCE THE BALL EXPERIENCES NO VERTICAL ACCELERATION, $T \cos \theta = Mg$ $M = \frac{T \cos \theta}{g}$

GO ON TO THE NEXT PAGE.

(c) Determine the speed of the ball.

THE NET FORCE ON THE BALL IS $T \sin \alpha$. HOWEVER, SINCE THE MOTION IS CIRCULAR, THE FORCE IS ALSO $\frac{mv^2}{r}$ ON, SINCE $r = l \sin \alpha$, $\frac{mv^2}{l \sin \alpha}$.

$$\text{So } T \sin \alpha = \frac{mv^2}{l \sin \alpha} ; v^2 = \frac{T l \sin^2 \alpha}{m} ; v^2 = \frac{T l \sin^2 \alpha}{\left(\frac{T \cos \alpha}{g}\right)} = g l \sin \alpha \tan \alpha.$$

$$v = \sqrt{g l \sin \alpha \tan \alpha} \text{ m/s}$$


(d) Determine the frequency of revolution of the ball.

$$\alpha = \frac{v}{r} = \frac{\sqrt{g l \sin \alpha \tan \alpha}}{l \sin \alpha} = \sqrt{\frac{g}{l \cos \alpha}}. \quad f = \frac{1}{2\pi} \alpha = \frac{1}{2\pi} \sqrt{\frac{g}{l \cos \alpha}} \text{ Hz}$$

(e) Suppose that the string breaks as the ball swings in its circular path. Qualitatively describe the trajectory of the ball after the string breaks but before it hits the ground.

IT WILL CONTINUE AT A CONSTANT SPEED IN THE COMPONENT OF VELOCITY TANGENT TO THE CURVE AT THE POINT OF BREAKAGE, BUT WILL EXPERIENCE A CONSTANT DOWNWARD ACCELERATION, AND SO WOULD CUT A PARABOLIC SHAPE THROUGH SPACE.


GO ON TO THE NEXT PAGE.



2. (15 points)

A ball attached to a string of length ℓ swings in a horizontal circle, as shown above, with a constant speed. The string makes an angle θ with the vertical, and T is the magnitude of the tension in the string. Express your answers to the following in terms of the given quantities and fundamental constants.

- (a) On the figure below, draw and label vectors to represent all the forces acting on the ball when it is at the position shown in the diagram. The lengths of the vectors should be consistent with the relative magnitudes of the forces.



- (b) Determine the mass of the ball.

$$T = \frac{2\pi r}{v}$$

$$m \frac{v^2}{r} = T \sin \theta$$

$$m \frac{v^2}{r \sin \theta} = T \sin \theta$$

$$m \frac{v^2}{r \sin \theta} = T \sin \theta$$

$$v^2 = \frac{r \ell \sin^2 \theta}{\cos \theta} g$$

$$T \cos \theta = mg$$

$$T = \frac{mg}{\cos \theta}$$

$$m = \frac{T \cos \theta}{g}$$

GO ON TO THE NEXT PAGE.

$$v^2 = g \frac{T}{2\pi} \tan \theta$$

(c) Determine the speed of the ball.

$$T = \frac{2\pi r}{v}$$

$$\frac{mv^2}{r} = T \sin \theta$$

$$v = \frac{g T \tan \theta}{2\pi}$$

$$r = \frac{T v}{2\pi}$$

$$m = \frac{T \cos \theta}{g}$$

$$\frac{\cancel{X} \cos \theta v^2}{g r} = \cancel{X} \sin \theta$$

$$v = \sqrt{\frac{g l \sin^2 \theta}{\cos \theta}}$$

$$r = l \sin \theta$$


$$v^2 = g r \tan \theta$$

(d) Determine the frequency of revolution of the ball.

$T = \frac{2\pi r}{v}$
 $f = \frac{v}{2\pi l \sin \theta}$
 $f = \left(\frac{g T \tan \theta}{2\pi} \right) \left(\frac{1}{2\pi l \sin \theta} \right)$
 $= \left(\frac{g T \tan \theta}{2\pi} \right) \left(\frac{1}{2\pi r} \right)$
 $f = \frac{\sqrt{\frac{g l \sin^2 \theta}{\cos \theta}}}{2\pi l \sin \theta} = \frac{\sin \theta \sqrt{\frac{g l}{\cos \theta}}}{2\pi l}$
 $= \frac{g T \tan \theta}{2\pi} \left(\frac{1}{2\pi r} \right) =$

(e) Suppose that the string breaks as the ball swings in its circular path. Qualitatively describe the trajectory of the ball after the string breaks but before it hits the ground.

At the moment the string broke, the ball would fly along a tangent from the circle it was originally traveling in. It would then follow a parabolic path to the ground.



GO ON TO THE NEXT PAGE.