

# Harold's Counting Cheat Sheet

22 October 2022

(See also Harold's Sets Cheat Sheet)

## Counting Rules

Rule	Description	Comments
<b>Cardinality</b>	The cardinality of A is the number of elements in set $A =  A $	<ul style="list-style-type: none"> <li>if <math>A = \{(1,2), (3,4), (5,6)\}</math>, then <math> A  = 3</math></li> <li>Also denoted <math>n(A)</math></li> <li>Cardinality = Counting</li> </ul>
<b>Product Rule</b>	Let $A_1, A_2, \dots, A_n$ be finite sets. Then, $ A_1 \times A_2 \times \dots \times A_n  =  A_1  \cdot  A_2  \cdot \dots \cdot  A_n $	<ul style="list-style-type: none"> <li>Counts sequences</li> <li>Think Intersection (<math>\cap</math>)</li> </ul>
<b>Sum Rule</b>	Consider n sets, $A_1, A_2, \dots, A_n$ . If the sets are mutually disjoint ( $A_i \cap A_j = \emptyset$ for $i \neq j$ ), then $ A_1 \cup A_2 \cup \dots \cup A_n  =  A_1  +  A_2  + \dots +  A_n $	<ul style="list-style-type: none"> <li>Counts sequences</li> <li>Think Union (<math>\cup</math>)</li> </ul>
<b>Generalized Product Rule</b>	$ S  = n_1 \cdot n_2 \cdot \dots \cdot n_k$ $n! = (n)(n-1)(n-2) \dots (2)(1)$	<ul style="list-style-type: none"> <li>In selecting an item from a set, if the number of choices at each step is independent, then the number of items in the set is the product of the number of choices in each step.</li> </ul>
<b>Bijection Rule</b>	Let S and T be two finite sets. If there is a bijection from S to T, then $ S  =  T $	<ul style="list-style-type: none"> <li>1-to-1 Correspondence</li> </ul>
<b>k-to-1 Rule</b>	$ Y  = \frac{ X }{k}$	<ul style="list-style-type: none"> <li>k-to-1 Correspondence</li> </ul>

## Counting Formulas & Techniques

Rule	Description	Comments
<b>Factorial</b>	$n!$ $= n \cdot (n - 1) \cdot (n - 2) \cdot \dots \cdot 3 \cdot 2 \cdot 1$	<ul style="list-style-type: none"> <li>The number of permutations of a finite set with n elements is P(n,n)</li> </ul>
<b>Permutation</b>	$P(n, r) = {}_n P_r = \frac{n!}{(n - r)!}$ $= n(n - 1) \dots (n - r + 1)$	<ul style="list-style-type: none"> <li>Order matters ( , )</li> <li>r-permutation</li> <li>Counting sequences</li> <li>Common application of the generalized product rule</li> <li>Order can be fixed but arbitrary</li> <li>Elements cannot be repeated</li> <li>Use when elements are all <u>different</u></li> </ul>
<b>Grouping</b>	Password length is 18. No character repeats. Must contain: a, z, 1, and 9.  $P(18, 4) \cdot P(36 - 4, 18 - 4)$	<ul style="list-style-type: none"> <li>Combines permutation with product rule</li> </ul>
<b>Combination</b>	$C(n, r) = {}_n C_r = \binom{n}{r}$ $= \frac{n!}{r! (n - r)!}$	<ul style="list-style-type: none"> <li>Order does not matter { , }</li> <li>Counting subsets</li> <li>r-combination</li> <li>n choose r</li> <li>Counting the r-subsets</li> <li>Combination = subset</li> <li>Use when elements are all <u>identical</u></li> </ul>
	Identity: $\binom{n}{n - r} = \binom{n}{r}$	<ul style="list-style-type: none"> <li>An equation is called an <b>identity</b> if the equation holds for all values for which the expressions in the equation are well defined.</li> </ul>
<b>Counting Subsets</b>	Bijection from: 5-bit strings with exactly 2 1's To: 2-subsets of { 1, 2, 3, 4, 5 } = $\binom{5}{2} = 10$	<ul style="list-style-type: none"> <li>Binary example</li> <li>Counting Strings by Counting Subsets</li> </ul>
	$\binom{m + n}{m}$	<ul style="list-style-type: none"> <li>Counting paths on a grid {N, E}</li> <li>Binary example if m = #1s &amp; n = #0s</li> </ul>

## Counting with Discrete Probability

Rule	Formula	Definition
<b>Notation</b>	$\cap$ = Intersection or "and" $\cup$ = Union or "or" $\bar{\quad}$ = Negation or "not"	<ul style="list-style-type: none"> <li>"and" implies multiplication.</li> <li>"or" implies addition.</li> <li>"not" implies negation.</li> </ul>
<b>Independent</b>	If $P(A B) = P(A)$	<ul style="list-style-type: none"> <li>The occurrence of one event does not affect the probability of the other event, or vice versa.</li> </ul>
<b>Mutually Independent</b>	No sets overlap. Future outcomes are not impacted by previous outcomes.	<ul style="list-style-type: none"> <li>Applies to more than two events</li> </ul>
<b>Dependent</b>	If $ A \cap B  \neq 0$	<ul style="list-style-type: none"> <li>The occurrence of one event affects the probability of the other event.</li> </ul>
<b>Disjoint</b> (“mutually exclusive”)	If $ A \cap B  = 0$ , then $ A \cup B  =  A  +  B $	<ul style="list-style-type: none"> <li>The events can never occur together.</li> </ul>
<b>Probability</b> (“likelihood”)	$P(E) = \frac{ E }{ S }$	<ul style="list-style-type: none"> <li><math>S</math> = Sample space or entire set</li> <li><math>A, B, E</math> = Event or subset</li> </ul> $0 \leq P(E) \leq 1$
<b>Addition Rule</b> (“or”)	$ A \cup B $ $=  A  +  B  -  A \cap B $	<b>Inclusion-Exclusion Principle</b> <ul style="list-style-type: none"> <li>Let <math>A, B</math> and <math>C</math> be three finite sets, then ...</li> <li>If sets overlap, then don't double count</li> <li>"... in any of the 3."</li> <li>"... divisible by 2, 3, or 5."</li> </ul>
	$ A \cup B \cup C $ $=  A  +  B  +  C  -  A \cap B  -  B \cap C  -  A \cap C  +  A \cap B \cap C $	
	$ A \cup B \cup C \cup D $ $=  A  +  B  +  C  +  D $ $-  A \cap B  -  A \cap C  -  A \cap D  -  B \cap C  -  B \cap D  -  C \cap D $ $+  A \cap B \cap C  +  A \cap B \cap D  +  A \cap C \cap D  +  B \cap C \cap D $ $-  A \cap B \cap C \cap D $	
	if mutually independent / disjoint: $ A_1 \cup A_2 \cup \dots \cup A_n  =  A_1  +  A_2  + \dots +  A_n $	<ul style="list-style-type: none"> <li>A collection of sets is <b>mutually disjoint</b> if the intersection of every pair of sets in the collection is empty.</li> <li>Restatement of the Sum Rule</li> </ul>

<p><b>Multiplication Rule</b> ("and")</p>	$ A \cap B  =  A  \cdot  (B   A) $ $ A \cap B  =  B  \cdot  (A   B) $ $ A \cap B  =  A  -  A \cap \bar{B} $ <p>if independent / disjoint:  <math display="block"> A \cap B  =  A  \cdot  B </math></p> <p>if mutually independent / disjoint:  <math display="block"> A \cap B \cap C  =  A  \cdot  B  \cdot  C </math> <math display="block"> A_1 \cap A_2 \cap \dots \cap A_n </math> <math display="block">=  A_1  \cdot  A_2  \cdot \dots \cdot  A_n </math></p>	
<p><b>Complement Rule / Subtraction Rule</b> ("not")</p>	$P(S) = P(E \cup \bar{E})$ $ E  +  \bar{E}  =  S $ $ E  =  S  -  \bar{E} $ $ \bar{E}  =  S  -  E $ $ (A   B)  +  (\bar{A}   B)  =  A $	<ul style="list-style-type: none"> <li>• S = entire set, E = subset</li> <li>• The complement of event E (denoted <math>\bar{E}</math> or <math>E^c</math>) means "not E";</li> <li>• It consists of all simple outcomes that are not in E.</li> <li>• "has at least one" so choose <math>\bar{E}</math> as "none"</li> </ul>
<p><b>Union by Compliment</b></p>	$ S  -  \overline{E_1 \cup E_2 \cup \dots \cup E_n}  =  E_1 \cup E_2 \cup \dots \cup E_n $	<ul style="list-style-type: none"> <li>• S = U = Universal set (all)</li> <li>• E.g., <math>10^4 - 9^4</math></li> </ul>
<p><b>Conditional Probability</b> ("given that")</p>	$P(A   B) = \frac{ A \cap B }{ B }$ <p>if independent / disjoint:  <math display="block">P(A   B) = \frac{P(A \cap B)}{P(B)} = P(A)</math> <math display="block"> (A   B)  =  A </math> <math display="block"> (B   A)  =  B </math></p>	<ul style="list-style-type: none"> <li>• Means the probability of event A given that event B has already occurred.</li> <li>• Is a rephrasing of the Multiplication Rule.</li> <li>• <math>P(A   B)</math> is the proportion of elements in B that are ALSO in A.</li> </ul>
<p><b>Total Probability Rule</b></p>	$P(A) = P(A \cap B_1) + \dots + P(A \cap B_n)$ $= P(B_1) \cdot P(A   B_1) + \dots + P(B_n) \cdot P(A   B_n)$ $P(A) = P(A \cap B) + P(A \cap \bar{B})$ $= P(A   B) \cdot P(B) + P(A   \bar{B}) \cdot P(\bar{B})$	<ul style="list-style-type: none"> <li>• To find the probability of event A, partition the sample space into several disjoint events.</li> <li>• A must occur along with one and only one of the disjoint events.</li> </ul>
<p><b>Bayes' Theorem</b></p>	$P(A   B) = \frac{ A \cap B }{ B } = \frac{ (B   A)  \cdot  A }{ B }$ $= \frac{ (B   A)  \cdot  A }{ (B   A)  \cdot  A  +  (B   \bar{A})  \cdot  \bar{A} }$	<ul style="list-style-type: none"> <li>• Allows <math>P(A   B)</math> to be calculated from <math>P(B   A)</math>.</li> <li>• Meaning it allows us to reverse the order of a conditional probability statement, and is the only generally valid method!</li> </ul>

**Sources:**

- [SNHU MAT 230](#) - Discrete Mathematics, zyBooks.