

# Harold's Ordinary Differential Equations (ODE) and Transforms Cheat Sheet

26 October 2026

## Classification

| Term                                        | Definition                                                                                                                                                                                                            |
|---------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>By Type</b>                              |                                                                                                                                                                                                                       |
| <b>Differential Equation (DiffEq)</b>       | A mathematical equation that relates a function to its derivatives.<br>Example:<br>$a_n(x) y^{(n)}(x) + a_{n-1}(x) y^{(n-1)}(x) + \cdots + a_1(x) y'(x) + a_0(x) y(x) = g(x)$                                         |
| <b>Ordinary Differential Equation (ODE)</b> | A differential equation that involves an unknown function and its derivatives with respect to a single independent variable.<br>Example:<br>$\frac{dy}{dx} = x^2 + 3x$                                                |
| <b>Partial Differential Equation (PDE)</b>  | A type of equation that involves an unknown function and its partial derivatives with respect to multiple independent variables.<br>Example:<br>$\frac{\partial u}{\partial t} + c \frac{\partial u}{\partial x} = 0$ |
| <b>Systems of Differential Equations</b>    | Two or more differential equations involving multiple unknown functions, solved together.<br>Example:<br>$\mathbf{x}' = A\mathbf{x}$                                                                                  |
| <b>By Order</b>                             |                                                                                                                                                                                                                       |
| <b>Order</b>                                | The order of the highest derivative in the equation.<br>$y' = 1^{\text{st}}$ order<br>$y'' = 2^{\text{nd}}$ order<br>$y''' = 3^{\text{rd}}$ order<br>$y^{(4)} = 4^{\text{th}}$ order                                  |
| <b>By Linearity</b>                         |                                                                                                                                                                                                                       |
| <b>Linear Equation</b>                      | $y^{(n)}(x) + a_{n-1}(x)y^{(n-1)}(x) + \cdots + a_1(x)y'(x) + a_0(x)y(x) = f(x)$ <i>where each <math>a_i(x)</math> is a function of <math>x</math>.</i>                                                               |
| <b>Nonlinear Equation</b>                   | An equation that contains functions of $y$ , such as $e^y$ , or functions of the derivatives of $y$ , such as $(5y)y''$ , $\sin(y')$ , or $\left(\frac{dy}{dx}\right)^2$ .                                            |
| <b>By Solution Type</b>                     |                                                                                                                                                                                                                       |
| <b>General Solution</b>                     | A family of functions that has parameters and does not take any initial conditions into account.                                                                                                                      |
| <b>Particular Solution</b>                  | A specific solution with no arbitrary constants. An IVP solution also satisfies the given initial conditions.                                                                                                         |
| <b>Singular Solution</b>                    | Cannot be obtained from the general solution.                                                                                                                                                                         |

## Notation and Terminology

| Notation                  | Expanded Form                                                                                                                                                | Description                                                                                          |
|---------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------|
| <b>Terms and Notation</b> |                                                                                                                                                              |                                                                                                      |
| $x, y, z, t$              | $x, y, z, t$                                                                                                                                                 | Variables for space and time.                                                                        |
| $f$                       | $f = f(x)$ or $y$                                                                                                                                            | The function of a single independent variable.                                                       |
| $f'$                      | $\frac{df}{dx}, \frac{dy}{dx}, \text{ or } y'$                                                                                                               | The derivative of $f$ with respect to $x$ .                                                          |
| $f_x$                     | $\frac{\partial f}{\partial x}$                                                                                                                              | The partial derivative of $f$ with respect to $x$ .<br>All other variables are treated as constants. |
| $y_p$                     | Particular solution to the equation.                                                                                                                         |                                                                                                      |
| <b>IVP</b>                | A differential equation together with initial conditions specified at the same point, so a particular solution can be found.<br>$y(0) = 1, \quad y'(0) = -1$ |                                                                                                      |



## Transforms

| Symbol                     | Name                               | Description                                                                                                                                                                                                                                                                                                          |
|----------------------------|------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| $\mathcal{L}\{f(t)\}$      | Laplace Transform                  | <p>Converts linear ODEs with constant coefficients into algebraic equations incorporating initial conditions. Especially useful for discontinuous or impulsive forcing.</p> $\mathcal{L}\{f(t)\} = F(s) = \int_0^{\infty} e^{-st} f(t) dt$                                                                           |
| $\mathcal{F}\{f(t)\}$      | Fourier Transform                  | <p>Transforms functions into the frequency domain. Useful for ODEs and PDEs on infinite or unbounded domains. Converts derivatives into algebraic factors.</p> $\mathcal{F}\{f(t)\} = \hat{f}(\omega) = \int_{-\infty}^{\infty} f(t) e^{-i\omega t} dt$                                                              |
| $F_s(k)$<br>&<br>$F_c(k)$  | Fourier Sine and Cosine Transforms | <p>Transform PDE boundary-value problems on <math>x \geq 0</math>. The sine transform naturally handles zero-value conditions at <math>x = 0</math>. The cosine transform naturally handles zero-slope conditions there.</p> $F_s(k) = \int_0^{\infty} f(x) \sin(kx) dx$ $F_c(k) = \int_0^{\infty} f(x) \cos(kx) dx$ |
| $\mathcal{H}_0\{f(r)\}(k)$ | Hankel Transform (Order 0)         | <p>Transforms radially symmetric PDEs on unbounded radial domains in cylindrical or circular geometry.</p> $\mathcal{H}_0\{f(r)\}(k) = F(k) = \int_0^{\infty} f(r) J_0(kr) r dr$ <p>where <math>J_0</math> is the Bessel function of the first kind, order zero.</p>                                                 |
| $\mathcal{M}\{f(x)\}(s)$   | Mellin Transform                   | <p>Transforms equations with scaling symmetry and certain singular or asymptotic problems.</p> $\mathcal{M}\{f(x)\}(s) = F(s) = \int_0^{\infty} x^{s-1} f(x) dx.$                                                                                                                                                    |
| $\mathcal{Z}\{x[n]\}$      | Z-Transform                        | <p>Transforms linear difference equations with constant coefficients into algebraic equations.</p> $\mathcal{Z}\{x[n]\} = X(z) = \sum_{n=0}^{\infty} x[n] z^{-n}$                                                                                                                                                    |

### Conditions

- Integral transforms require convergence on their stated domains.

## Direction Fields

| Name                                    | Slope Equation                                                                                                             | Graph                                                                               |
|-----------------------------------------|----------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------|
| <b>Direction Field</b><br>(Slope Field) | $y' = x + y$<br><b>General Solution:</b><br>$y(x) = c_1 e^x - x - 1$                                                       |   |
| <b>Solution Curve</b>                   | $\frac{dy}{dx} = x + y$<br><b>Initial Condition:</b><br>$y(0) = 1$<br><b>Particular Solution:</b><br>$y(x) = 2e^x - x - 1$ |  |

# 1<sup>st</sup>-Order Methods

| Technique                                               | Form of Equation                                                                                                                                                                                                          | Solution Method                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
|---------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>1<sup>st</sup>-Order Linear</b>                      | $y' + a(x)y = f(x)$                                                                                                                                                                                                       | $I.F. = e^{\int a(x) dx}$<br>$y = e^{-A(x)} \left[ C + \int e^{A(x)} f(x) dx \right]$<br>$A(x) = \int a(x) dx$                                                                                                                                                                                                                                                                                                                                                                                                      |
| <b>Single Variable</b>                                  | $\frac{dy}{dx} = f(x)$                                                                                                                                                                                                    | $y(x) = \int f(x) dx + C$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| <b>Separable Equations</b><br>(Separation of Variables) | $\frac{dy}{dx} = f(x, y) = \frac{g(x)}{f(y)}$                                                                                                                                                                             | $\int f(y) dy = \int g(x) dx + C$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| <b>Integrating Factors</b><br>$(\mu(x))$                | $\frac{dy}{dx} + P(x)y = Q(x)$                                                                                                                                                                                            | <ol style="list-style-type: none"> <li><math>I.F. = \mu(x) = e^{\int P(x) dx}</math></li> <li>Multiply the entire equation by <math>\mu(x)</math>.<br/> <math>\mu(x) \frac{dy}{dx} + \mu(x)P(x)y = \mu(x)Q(x)</math></li> <li>But the derivative of the product is:<br/> <math>\frac{d(\mu(x)y(x))}{dx} = \mu(x) \frac{dy}{dx} + \mu(x)P(x)y</math></li> <li><math>\mu(x)y(x) = \int \mu(x) Q(x) dx + C</math></li> <li>General solution<br/> <math>y(x) = \frac{\int \mu(x) Q(x) dx + C}{\mu(x)}</math></li> </ol> |
| <b>Exact Equations</b>                                  | $M(x, y) dx + N(x, y) dy = 0$<br>$= d\Phi(x, y)$<br><br>$If f \quad \frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$<br><br>Assume $M, N$ have continuous partial derivatives on a simply connected domain. | Find a potential function $\Phi(x, y)$ such that<br>$\Phi_x = M, \quad \Phi_y = N.$<br><br>The implicit solution is $\Phi(x, y) = C.$<br>Find it by integrating and comparing:<br>$\int M(x, y) dx \quad and \quad \int N(x, y) dy$                                                                                                                                                                                                                                                                                 |
| <b>Non-Exact Equations</b>                              | Reduction to Exact via Integrating Factor.                                                                                                                                                                                |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
|                                                         | $\mu(x, y) \left[ M(x, y) + N(x, y) \frac{dy}{dx} \right] = 0$                                                                                                                                                            |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
|                                                         | <b>If</b>                                                                                                                                                                                                                 | <b>Then</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
|                                                         | <b>Case 1: <math>\mu(x)</math></b><br>$h_1(x) \equiv \frac{M_y - N_x}{N}$                                                                                                                                                 | $\mu(x, y) = e^{\int h_1(x) dx}$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
|                                                         | <b>Case 2: <math>\mu(y)</math></b><br>$h_2(y) \equiv \frac{N_x - M_y}{M}$                                                                                                                                                 | $\mu(x, y) = e^{\int h_2(y) dy}$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
|                                                         | <b>Case 3: <math>\mu(xy)</math></b><br>$h_3(xy) \equiv \frac{N_x - M_y}{xM - yN}$                                                                                                                                         | $\mu(x, y) = e^{\int h_3(xy) d(xy)}$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |

|                                                 |                                                                                                                                                                                                                              |                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
|-------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Bernoulli</b>                                | $\frac{dy}{dx} + P(x)y = Q(x)y^n$                                                                                                                                                                                            | <p>If <math>n = 0</math> or <math>n = 1</math>, this is linear; else</p> <ol style="list-style-type: none"> <li>1. Divide by <math>y^n</math></li> <li>2. Change variable <math>z = y^{1-n}</math></li> <li>3. Apply substitution to get a linear form</li> <li>4. Use an integrating factor to solve</li> <li>5. Check separately for solutions excluded by division or substitution, including <math>y = 0</math> when applicable.</li> </ol>     |
| <b>Homogeneous Substitution</b><br>( $y = vx$ ) | $M(x, y) dx + N(x, y) dy = 0$<br>where $M(x, y)$ and $N(x, y)$ are homogeneous functions of the same degree $n$ , meaning:<br>$M(\lambda x, \lambda y) = \lambda^n M(x, y)$<br>$N(\lambda x, \lambda y) = \lambda^n N(x, y)$ | <ol style="list-style-type: none"> <li>1. Start with:<br/> <math display="block">\frac{dy}{dx} = F\left(\frac{y}{x}\right)</math></li> <li>2. Reduce to separation of variables using:<br/> <math display="block">v = \frac{y}{x}</math> <math display="block">y = vx</math></li> <li>3. Product rule:<br/> <math display="block">\frac{dy}{dx} = v + x \frac{dv}{dx}</math></li> <li>4. Rewrite in terms of <math>\frac{dv}{dx}</math>.</li> </ol> |
| <b>Substitution</b>                             | $\frac{dy}{dx} = F(ax + by + c)$<br>$b \neq 0$                                                                                                                                                                               | <ol style="list-style-type: none"> <li>1. Set <math>v = ax + by + c</math></li> <li>2. Solve for <math>y</math></li> <li>3. Find <math>\frac{dy}{dx}</math> in terms of <math>\frac{dv}{dx}</math></li> <li>4. Rewrite the equation in terms of <math>\frac{dv}{dx}</math></li> </ol>                                                                                                                                                               |
| <b>Reduction by Translation</b>                 | $y' = \frac{Ax + By + C}{Dx + Ey + F}$                                                                                                                                                                                       |                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
|                                                 | <b>If</b>                                                                                                                                                                                                                    | <b>Then</b>                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| <b>Case 1:<br/>Intersecting Lines</b>           | $AE - BD \neq 0$                                                                                                                                                                                                             | <ol style="list-style-type: none"> <li>1. Set <math>x = X + h</math>, <math>y = Y + k</math></li> <li>2. Solve the system of equations:<br/> <math display="block">Ah + Bk + C = 0</math> <math display="block">Dh + Ek + F = 0</math></li> <li>3. Use <math>Y = vX</math></li> <li>4. Solve using separation of variables</li> <li>5. Translate back</li> </ol>                                                                                    |
| <b>Case 2:<br/>Parallel Lines</b>               | $AE - BD = 0$<br>$B \neq 0$                                                                                                                                                                                                  | <ol style="list-style-type: none"> <li>1. Put<br/> <math display="block">u = Ax + By</math> <math display="block">y' = \frac{u' - A}{B}</math></li> <li>2. Solve</li> </ol>                                                                                                                                                                                                                                                                         |

## 2<sup>nd</sup>-Order Linear Homogeneous (= 0)

| Technique                                                 | Form of Equation                                                                                                                                                                                         | Solution Method                                                                                                                                                                                                                                                                                                  |
|-----------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>2<sup>nd</sup>-Order Linear</b>                        | $P(x)y'' + Q(x)y' + R(x)y = 0$                                                                                                                                                                           | The typical case has all constant ( $a, b, c$ ) coefficients instead of functions of $x$ .                                                                                                                                                                                                                       |
| <b>Homogeneous</b>                                        | $ay'' + by' + cy = 0$                                                                                                                                                                                    | $y(x) = c_1y_1(x) + c_2y_2(x)$<br>where $y_1, y_2$ are linearly independent solutions.                                                                                                                                                                                                                           |
| <b>Principle of Superposition</b>                         | $y'' + ay' + by = L[y]$                                                                                                                                                                                  | <b>Case 1:</b><br>If $L[y_1] = 0$ and $L[y_2] = 0$ , then<br>$y = c_1y_1 + c_2y_2$<br><b>Case 2:</b><br>If $L[y_1] = f_1$ and $L[y_2] = f_2$ , then<br>$L[y_1 + y_2] = f_1 + f_2$                                                                                                                                |
| <b>Characteristic Equation</b><br>(Constant Coefficients) | $ay'' + by' + cy = 0$<br>$\Downarrow$<br>$ar^2 + br + c = 0$                                                                                                                                             | Try a solution of the form:<br>$y(x) = e^{rx}$<br>Roots: $r_1, r_2$                                                                                                                                                                                                                                              |
|                                                           | <b>If</b>                                                                                                                                                                                                | <b>Then</b>                                                                                                                                                                                                                                                                                                      |
| <b>Case 1:<br/>Distinct Roots</b>                         | $r_1, r_2 \in \mathbb{R}$<br>$r_1 \neq r_2$                                                                                                                                                              | 1. Set<br>$y_1(x) = e^{r_1x}$<br>$y_2(x) = e^{r_2x}$<br>2. Superposition gives<br>$y(x) = c_1e^{r_1x} + c_2e^{r_2x}$                                                                                                                                                                                             |
| <b>Case 2:<br/>Repeated Root</b>                          | $r_1, r_2 \in \mathbb{R}$<br>$r_1 = r_2 = r$                                                                                                                                                             | $y(x) = c_1e^{rx} + c_2xe^{rx}$                                                                                                                                                                                                                                                                                  |
| <b>Case 3:<br/>Complex Roots</b>                          | $r_{1,2} = \lambda \pm \mu i$<br>with<br>$\lambda = -\frac{b}{2a}, \quad \mu = \frac{\sqrt{4ac - b^2}}{2a}$                                                                                              | 1. Set<br>$y_1(x) = e^{(\lambda + \mu i)x}$<br>$y_2(x) = e^{(\lambda - \mu i)x}$<br>2. Superposition gives<br>$y(x) = c_1e^{(\lambda + \mu i)x} + c_2e^{(\lambda - \mu i)x}$<br>3. Apply $e^{i\theta} = \cos(\theta) + i\sin(\theta)$<br>4. $y(x) = c_1e^{\lambda x} \cos(\mu x) + c_2e^{\lambda x} \sin(\mu x)$ |
| <b>Reduction of Order</b>                                 | $P(x)y'' + Q(x)y' + R(x)y = 0$                                                                                                                                                                           | If we already know $y_1$ ,<br>1. Put $y_2 = vy_1$<br>2. Expand in terms of $v'', v', v$<br>3. Put $z = v'$<br>4. Solve the reduced equation                                                                                                                                                                      |
| <b>Wronskian</b><br>(Linear Independence)                 | For two solutions of the same regular 2 <sup>nd</sup> -order linear homogeneous ODE $[y_1(x) \text{ and } y_2(x)]$ , linear independence is equivalent to $W(x_0) \neq 0$ at some point in the interval. | $W(y_1, y_2)(x) = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} \neq 0$                                                                                                                                                                                                                                 |

|                                   |                                                                                                                                  |                                                                                                                                                             |
|-----------------------------------|----------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Euler-Cauchy Equation</b>      | $x^2 y'' + axy' + by = 0$ <p>where <math>x &gt; 0</math></p> <p>A. E. : <math>\lambda(\lambda - 1) + a\lambda + b = 0</math></p> | 1. $y(x)$ of the form $x^\lambda$<br>2. Reduction to Constant Coefficients:<br>Use $x = e^t, t = \ln x$<br>3. Rewrite in terms of $t$ using the chain rule. |
| <b>Case 1:<br/>Distinct Roots</b> | $\lambda_1, \lambda_2 \in \mathbb{R}$<br>$\lambda_1 \neq \lambda_2$<br>$x > 0$                                                   | $y(x) = c_1 x^{\lambda_1} + c_2 x^{\lambda_2}$                                                                                                              |
| <b>Case 2:<br/>Repeated Root</b>  | $\lambda_1, \lambda_2 \in \mathbb{R}$<br>$\lambda_1 = \lambda_2 = \lambda$<br>$x > 0$                                            | $y(x) = x^\lambda (c_1 + c_2 \ln x )$                                                                                                                       |
| <b>Case 3:<br/>Complex Roots</b>  | $\lambda_{1,2} = \alpha \pm i\beta$<br>$x > 0$                                                                                   | $y(x) = x^\alpha (c_1 \cos(\beta \ln x )$<br>$+ c_2 \sin(\beta \ln x ))$                                                                                    |

## 2<sup>nd</sup>-Order Linear Non-Homogeneous: Variation of Parameters (= $g(x)$ )

| Technique                          | Form of Equation              | Solution Method                                                                                   |
|------------------------------------|-------------------------------|---------------------------------------------------------------------------------------------------|
| <b>2<sup>nd</sup>-Order Linear</b> | $y'' + P(x)y' + Q(x)y = g(x)$ | Works beyond constant-coefficient equations when the homogeneous fundamental solutions are known. |
| <b>Variation of Parameters</b>     | $y'' + P(x)y' + Q(x)y = g(x)$ | $y_p = -y_1 \int \frac{y_2 g}{W} dx + y_2 \int \frac{y_1 g}{W} dx$ $W = y_1 y_2' - y_1' y_2$      |
| <b>General Solution</b>            |                               | $y = c_1 y_1 + c_2 y_2 + y_p$                                                                     |

### Conditions

- $y_1, y_2$  are linearly independent solutions of the associated homogeneous equation.
- $W \neq 0$  on the working interval.

## 2<sup>nd</sup>-Order Methods

| Technique                                                | Form of Equation                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                | Solution Method                                                                                                                                                                                                                                                                 |
|----------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>2<sup>nd</sup>-Order Linear</b>                       | $y'' + a(x)y' + b(x)y = f(x)$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   | $y = y_h + y_p$ $= c_1 y_1(x) + c_2 y_2(x) + y_p(x)$ <p>where <math>y_1, y_2</math> are linearly independent solutions, and the general solution includes the particular solution, <math>y_p</math>.</p>                                                                        |
| <b>Non-Homogeneous*</b>                                  | $P(x)y'' + Q(x)y' + R(x)y = f(x)$ $ay'' + by' + cy = f(x)$ $f(x) \not\equiv 0$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  | If homogeneous, then $f(x) = 0$ .                                                                                                                                                                                                                                               |
| <b>Case 1: <math>y', y</math> Missing</b>                | $y'' = f(x)$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    | $y(x) = \int \left( \int f(x) dx \right) dx + C_1 x + C_2$                                                                                                                                                                                                                      |
| <b>Case 2: <math>y</math> Missing</b>                    | $y'' = f(y', x)$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                | <ol style="list-style-type: none"> <li>1. Change of variable: <math>p = y'</math></li> <li>2. Solve the 1<sup>st</sup>-order equation for <math>p</math>.</li> <li>3. Then solve <math>y' = p</math></li> </ol>                                                                 |
| <b>Case 3: <math>x</math> Missing</b>                    | $y'' = f(y', y)$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                | <ol style="list-style-type: none"> <li>1. Change of variable: <math>p = y' +</math> chain rule</li> <li>2. <math>p \frac{dp}{dy} = f(p, y)</math> is a 1<sup>st</sup>-order ODE</li> <li>3. Solve it</li> <li>4. Back-replace <math>p</math></li> <li>5. Solve again</li> </ol> |
| <b>Case 4: <math>y', x</math> Missing</b>                | $y'' = f(y)$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    | <ol style="list-style-type: none"> <li>1. Change of variable: <math>p = y' +</math> chain rule</li> <li>2. <math>p \frac{dp}{dy} = f(y)</math> is a separation of vars</li> <li>3. Solve it</li> <li>4. Back-substitute <math>p</math></li> <li>5. Solve again</li> </ol>       |
| <b>Undetermined Coefficients</b><br>(Educated Guesswork) | <p>Choose a trial particular solution <math>y_p(x)</math> based on the form of <math>f(x)</math>.<br/>Valid forms:</p> <ol style="list-style-type: none"> <li>1. <math>P_n(x)</math></li> <li>2. <math>P_n(x) e^{ax}</math></li> <li>3. <math>e^{ax} (P_n(x) \cos bx + Q_n(x) \sin bx)</math></li> </ol> <p><u>Procedure:</u></p> <ul style="list-style-type: none"> <li>• Choose a trial solution containing all terms of the appropriate polynomial, exponential, sine, or cosine form.</li> <li>• For each trial group that overlaps the homogeneous solution, multiply that group by <math>x^m</math>, where <math>m</math> is the multiplicity of the corresponding characteristic root.</li> <li>• Substitute into the ODE and solve for the unknown coefficients.</li> </ul> <p>NOTE: This method applies to linear ODEs with constant coefficients.</p> |                                                                                                                                                                                                                                                                                 |

\*Cases 2-4 can be nonlinear.

## Laplace Transforms

| Technique                        | Equations and Descriptions                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
|----------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Laplace Transform                | $\mathcal{L}\{f(t)\} = F(s)$ <ul style="list-style-type: none"> <li>An integral transform that converts a function of a real variable (usually <math>t</math>, in the time domain) to a function of a complex variable <math>s</math> in the complex-valued frequency domain (<math>s</math> domain). <ul style="list-style-type: none"> <li>NOTE: Frequency <math>s = \sigma + i\omega</math> is a complex transform variable with units of inverse time.</li> </ul> </li> <li>Transforms ODEs into algebraic equations.</li> <li>The z-transform is the discrete analog of the Laplace transform in signal processing.</li> </ul> |
| Inverse Laplace Transform        | $\mathcal{L}^{-1}\{F(s)\} = f(t)$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| Laplace Transform Equation       | $\mathcal{L}\{f(t)\} = F(s) = \int_0^{\infty} e^{-st} f(t) dt$ <p>where</p> <ul style="list-style-type: none"> <li><math>f(t)</math> must be defined on <math>0 \leq t \leq \infty</math></li> <li><math>s</math> lies in the region of convergence; for real <math>s</math>, take <math>s</math> sufficiently large.</li> <li>The improper integral must converge</li> </ul>                                                                                                                                                                                                                                                       |
| Transform of a Derivative        | <b>Theorem:</b> If $f(t), f'(t), \dots, f^{(n-1)}(t)$ are continuous on $[0, \infty)$ and are of exponential order, and if $f^{(n)}(t)$ is piecewise continuous on $[0, \infty)$ , then the Laplace transform of the $n^{\text{th}}$ derivative of $f(t)$ is given by the general $n^{\text{th}}$ order derivative equation below.                                                                                                                                                                                                                                                                                                  |
| Laplace Transform of Derivatives | 1 <sup>st</sup> order derivative $\mathcal{L}\{f'(t)\} = s \mathcal{L}\{f(t)\} - f(0)$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
|                                  | 2 <sup>nd</sup> order derivative $\mathcal{L}\{f''(t)\} = s^2 \mathcal{L}\{f(t)\} - s f(0) - f'(0)$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
|                                  | 3 <sup>rd</sup> order derivative $\mathcal{L}\{f'''(t)\} = s^3 \mathcal{L}\{f(t)\} - s^2 f(0) - s f'(0) - f''(0)$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
|                                  | $n^{\text{th}}$ order derivative $\mathcal{L}\{f^{(n)}(t)\} = s^n \mathcal{L}\{f(t)\} - s^{n-1} f(0) - s^{n-2} f'(0) - \dots - f^{(n-1)}(0)$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| Using Laplace to Solve a DiffEq  | Solves IVPs for linear differential equations with constant coefficients. <ol style="list-style-type: none"> <li>Apply the Laplace transform to each term in the IVP.</li> <li>Solve the resulting algebraic equation for the transformed function <math>F(s)</math>.</li> <li>Apply the inverse Laplace transform to find the solution.</li> </ol>                                                                                                                                                                                                                                                                                 |

## Table of Laplace Transforms

| #  | $f(t)$                          | $\mathcal{L}\{f(t)\} = F(s)$                                                  | Restrictions                    |
|----|---------------------------------|-------------------------------------------------------------------------------|---------------------------------|
| 1  | $f'(t)$                         | $sF(s) - f(0)$                                                                |                                 |
| 2  | $f''(t)$                        | $s^2F(s) - sf(0) - f'(0)$                                                     |                                 |
| 3  | $f'''(t)$                       | $s^3F(s) - s^2f(0) - sf'(0) - f''(0)$                                         |                                 |
| 4  | $f^{(4)}(t)$                    | $s^4F(s) - s^3f(0) - s^2f'(0) - sf''(0) - f'''(0)$                            |                                 |
| 5  | $f^{(n)}(t)$                    | $s^nF(s) - s^{n-1}f(0) - s^{n-2}f'(0) - \dots - sf^{(n-2)}(0) - f^{(n-1)}(0)$ |                                 |
| 6  | 1                               | $\frac{1}{s}$                                                                 |                                 |
| 7  | $e^{at}$                        | $\frac{1}{s-a}, s > a$                                                        | $s > a$                         |
| 8  | $te^{at}$                       | $\frac{1}{(s-a)^2}$                                                           | $s > a$                         |
| 9  | $t^n e^{at}$                    | $\frac{n!}{(s-a)^{n+1}}$                                                      | $s > a$<br>$n = 1, 2, 3, \dots$ |
| 10 | $\frac{e^{at} - e^{bt}}{a-b}$   | $\frac{1}{(s-a)(s-b)}$                                                        | $a \neq b$<br>$s > \max(a, b)$  |
| 11 | $\frac{ae^{at} - be^{bt}}{a-b}$ | $\frac{s}{(s-a)(s-b)}$                                                        | $a \neq b$<br>$s > \max(a, b)$  |
| 12 | $t$                             | $\frac{1}{s^2}$                                                               |                                 |
| 13 | $t^n$                           | $\frac{n!}{s^{n+1}}$                                                          | $n = 1, 2, 3, \dots$            |
| 14 | $t^p$                           | $\frac{\Gamma(p+1)}{s^{p+1}}$                                                 | $p \in \mathbb{R}, p > -1$      |
| 15 | $\sqrt{t}$                      | $\frac{\sqrt{\pi}}{2s^{3/2}}$                                                 |                                 |
| 16 | $t^{n-\frac{1}{2}}$             | $\frac{1 \cdot 3 \cdot 5 \dots (2n-1)\sqrt{\pi}}{2^n s^{n+\frac{1}{2}}}$      | $n = 1, 2, 3, \dots$            |
| 17 | $\sin(at)$                      | $\frac{a}{s^2 + a^2}$                                                         |                                 |
| 18 | $\cos(at)$                      | $\frac{s}{s^2 + a^2}$                                                         |                                 |
| 19 | $\sin(at+b)$                    | $\frac{s \sin(b) + a \cos(b)}{s^2 + a^2}$                                     |                                 |
| 20 | $\cos(at+b)$                    | $\frac{s \cos(b) - a \sin(b)}{s^2 + a^2}$                                     |                                 |
| 21 | $t \sin(at)$                    | $\frac{2as}{(s^2 + a^2)^2}$                                                   |                                 |
| 22 | $t \cos(at)$                    | $\frac{s^2 - a^2}{(s^2 + a^2)^2}$                                             |                                 |
| 23 | $\frac{1}{t} \sin(at)$          | $\arctan\left(\frac{a}{s}\right)$                                             |                                 |
| 24 | $e^{at} \sin(bt)$               | $\frac{b}{(s-a)^2 + b^2}$                                                     | $s > a$                         |


|    |                                                         |                                                                                                            |                                                                                                                          |
|----|---------------------------------------------------------|------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------|
| 25 | $e^{at} \cos(bt)$                                       | $\frac{s-a}{(s-a)^2 + b^2}$                                                                                | $s > a$                                                                                                                  |
| 26 | $\sin(at) - at \cos(at)$                                | $\frac{2a^3}{(s^2 + a^2)^2}$                                                                               |                                                                                                                          |
| 27 | $\sin(at) + at \cos(at)$                                | $\frac{2as^2}{(s^2 + a^2)^2}$                                                                              |                                                                                                                          |
| 28 | $\cos(at) - at \sin(at)$                                | $\frac{s(s^2 - a^2)}{(s^2 + a^2)^2}$                                                                       |                                                                                                                          |
| 29 | $\cos(at) + at \sin(at)$                                | $\frac{s(s^2 + 3a^2)}{(s^2 + a^2)^2}$                                                                      |                                                                                                                          |
| 30 | $\sinh(at)$                                             | $\frac{a}{s^2 - a^2}$                                                                                      | $s >  a $                                                                                                                |
| 31 | $\cosh(at)$                                             | $\frac{s}{s^2 - a^2}, s >  a $                                                                             | $s >  a $                                                                                                                |
| 32 | $t \sinh(at)$                                           | $\frac{2as}{(s^2 - a^2)^2}$                                                                                | $s >  a $                                                                                                                |
| 33 | $t \cosh(at)$                                           | $\frac{s^2 + a^2}{(s^2 - a^2)^2}$                                                                          | $s >  a $                                                                                                                |
| 34 | $e^{at} \sinh(bt)$                                      | $\frac{b}{(s-a)^2 - b^2}$                                                                                  | $s > a +  b $                                                                                                            |
| 35 | $e^{at} \cosh(bt)$                                      | $\frac{s-a}{(s-a)^2 - b^2}$                                                                                | $s > a +  b $                                                                                                            |
| 36 | $e^{at} f(t)$                                           | $F(s-a)$                                                                                                   |                                                                                                                          |
| 37 | $tf(t)$                                                 | $-F'(s)$                                                                                                   |                                                                                                                          |
| 38 | $t^n f(t)$                                              | $(-1)^n F^{(n)}(s)$                                                                                        | $n = 1, 2, 3, \dots$                                                                                                     |
| 39 | $\frac{1}{t} f(t)$                                      | $\int_s^\infty F(u) du$<br>(Requires convergence of both sides,<br>including integrability near $t = 0$ .) |                                                                                                                          |
| 40 | $f(ct)$                                                 | $\frac{1}{c} F\left(\frac{s}{c}\right), c > 0$                                                             |                                                                                                                          |
| 41 | $u_a(t) = u(t-a)$<br><a href="#">Heaviside Function</a> | $\frac{1}{s} e^{-as}$                                                                                      | Nonnegative delay<br>$a \geq 0$                                                                                          |
| 42 | $\delta(t)$                                             | 1                                                                                                          |                                                                                                                          |
| 43 | $\delta(t-a)$<br><a href="#">Dirac Delta Function</a>   | $e^{-as}$                                                                                                  | Nonnegative delay<br>$a \geq 0$ .<br>(Using $a > 0$ for<br>the shifted<br>impulse avoids<br>the endpoint<br>convention.) |
| 44 | $u_a(t) f(t-a)$                                         | $e^{-as} F(s)$                                                                                             |                                                                                                                          |
| 45 | $u_a(t) g(t)$                                           | $e^{-as} \mathcal{L}\{g(t+a)\}$                                                                            |                                                                                                                          |
| 46 | $\int_0^t f(v) dv$                                      | $\frac{1}{s} F(s)$                                                                                         |                                                                                                                          |
| 47 | $\int_0^t f(t-\tau) g(\tau) d\tau$                      | $F(s) G(s)$                                                                                                |                                                                                                                          |

|    |                                                       |                                                |                                                                 |
|----|-------------------------------------------------------|------------------------------------------------|-----------------------------------------------------------------|
| 48 | $f(t + T) = f(t)$                                     | $\frac{\int_0^T e^{-st} f(t) dt}{1 - e^{-sT}}$ | $T > 0$ , with the periodic function integrable over one period |
| 49 | $af(t) + bg(t)$                                       | $aF(s) + bG(s)$                                |                                                                 |
| 50 | $\frac{1}{\sqrt{\pi t}} e^{\frac{-a^2}{4t}}$          | $\frac{e^{-a\sqrt{s}}}{\sqrt{s}}$              | $a > 0$                                                         |
| 51 | $\frac{a}{2\sqrt{\pi t^3}} e^{\frac{-a^2}{4t}}$       | $e^{-a\sqrt{s}}$                               | $a > 0$                                                         |
| 52 | $\operatorname{erfc}\left(\frac{a}{2\sqrt{t}}\right)$ | $\frac{e^{-a\sqrt{s}}}{s}$                     | $a > 0$                                                         |

### Conditions

- Valid in the region of convergence, typically  $s > 0$ .
- $\Gamma$  is the Gamma function, which extends factorials beyond nonnegative integers. For  $n = 0, 1, 2, \dots$ ,  $\Gamma(n + 1) = n!$ .

## Fourier Transforms

| Technique                                  | Equations and Descriptions                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
|--------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Fourier Series                             | <ul style="list-style-type: none"> <li>A method to <b>decompose</b> a periodic signal into a sum of sine and cosine functions with different frequencies and amplitudes.</li> <li>Is particularly useful for analyzing arbitrary <b>periodic</b> signals.</li> <li>Was introduced by Joseph Fourier.</li> <li>Is widely used in fields such as electronics, signal processing, and quantum mechanics.</li> </ul>                                                                                                                                                                           |
| Fourier Transform                          | <ul style="list-style-type: none"> <li>A mathematical operation that <b>transforms</b> a signal from the time domain (<math>t</math>) to the frequency domain (<math>\omega</math>).</li> <li>Unlike the Fourier Series, the Fourier Transform can be applied to both <b>periodic</b> and <b>non-periodic</b> signals.</li> <li>Is widely used to solve differential equations.</li> </ul>                                                                                                               |
| Fourier Transform Equation                 | $\mathcal{F}\{f(t)\} = \hat{f}(\omega) = \int_{-\infty}^{\infty} f(t) e^{-i\omega t} dt$ <p>where</p> <ul style="list-style-type: none"> <li><math>f(t)</math> must be defined on <math>-\infty \leq t \leq \infty</math></li> <li><math>\omega</math> is an arbitrary real variable where <math>\omega = 2\pi f</math></li> <li>The improper integral must converge</li> </ul>                                                                                                                                                                                                            |
| Inverse Fourier Transform Equation         | $\mathcal{F}^{-1}\{\hat{f}(\omega)\} = f(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \hat{f}(\omega) e^{i\omega t} d\omega$ <p>where</p> <ul style="list-style-type: none"> <li><math>\hat{f}(\omega)</math> must be defined on <math>-\infty \leq \omega \leq \infty</math></li> <li><math>\omega</math> is an arbitrary real variable where <math>\omega = 2\pi f</math></li> <li>The improper integral must converge</li> </ul> <p>Under standard inversion conditions, the inverse gives <math>f(t)</math> at continuity points and the average of the one-sided limits at a jump.</p> |
| Using Fourier Transforms to Solve a DiffEq | <p>Solves ODEs for linear differential equations with constant coefficients.</p> <ol style="list-style-type: none"> <li>1. Apply the Fourier transform to each term in the ODE.</li> <li>2. Solve the resulting algebraic equation for the transformed function <math>\hat{f}(\omega)</math>.</li> <li>3. Apply the inverse Fourier transform to find the solution.</li> </ol>                                                                                                                                                                                                             |
| Fourier Qualifications                     | <ul style="list-style-type: none"> <li>Ordinary transforms use convergent integrals.</li> <li>Entries involving constants, sinusoids, polynomials, or delta functions are interpreted as distributions.</li> <li>For <math>1/x</math>, use principal value; for <math>1/x^n, n \geq 2</math>, use finite part.</li> </ul>                                                                                                                                                                                                                                                                  |

|                |                                                                                                                                                                                                                                                           |
|----------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Example</b> | $f(t) = \begin{cases} e^{-at} & \text{for } t > 0 \\ 0 & \text{for } t < 0 \end{cases}$ $a > 0 \text{ for real } a$ $\hat{f}(\omega) = \int_{-\infty}^{\infty} f(t) e^{-i\omega t} dt = \int_0^{\infty} e^{-at} e^{-it\omega} dt = \frac{1}{a + i\omega}$ |
|----------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

**Table of Fourier Transforms ( $a, b > 0$ )**

| #  | $f(x)$                    | $\mathcal{F}\{f(x)\} = \hat{f}(\omega)$                                             |
|----|---------------------------|-------------------------------------------------------------------------------------|
| 1  | $a f(x) + b g(x)$         | $a \hat{f}(\omega) + b \hat{g}(\omega)$                                             |
| 2  | $f(x - a)$                | $e^{-ia\omega} \hat{f}(\omega)$                                                     |
| 3  | $f(x) e^{iax}$            | $\hat{f}(\omega - a)$                                                               |
| 4  | $f(ax)$                   | $\frac{1}{ a } \hat{f}\left(\frac{\omega}{a}\right)$                                |
| 5  | $\frac{d^n f(x)}{dx^n}$   | $(i\omega)^n \hat{f}(\omega)$                                                       |
| 6  | $x^n f(x)$                | $i^n \frac{d^n \hat{f}(\omega)}{d\omega^n}$                                         |
| 7  | $(f * g)(x)$              | $\hat{f}(\omega) \hat{g}(\omega)$                                                   |
| 8  | $f(x) g(x)$               | $\frac{1}{2\pi} (\hat{f} * \hat{g})(\omega)$                                        |
| 9  | Real $f(x)$               | $\hat{f}(-\omega) = \overline{\hat{f}(\omega)}$                                     |
| 10 | Imaginary $f(x)$          | $\hat{f}(-\omega) = -\overline{\hat{f}(\omega)}$                                    |
| 11 | $\overline{f(x)}$         | $\overline{\hat{f}(-\omega)}$                                                       |
| 12 | $f(x) \cos(ax)$           | $\frac{\hat{f}(\omega - a) + \hat{f}(\omega + a)}{2}$                               |
| 13 | $f(x) \sin(ax)$           | $\frac{\hat{f}(\omega - a) - \hat{f}(\omega + a)}{2i}$                              |
| 14 | $e^{-ax} u(x)$            | $\frac{1}{a + i\omega}$                                                             |
| 15 | $e^{-a x }$               | $\frac{2a}{a^2 + \omega^2}$                                                         |
| 16 | 1                         | $2\pi\delta(\omega)$                                                                |
| 17 | $\delta(x)$               | 1                                                                                   |
| 18 | $\sin(ax)$                | $-i\pi(\delta(\omega - a) - \delta(\omega + a))$                                    |
| 19 | $\cos(ax)$                | $\pi(\delta(\omega - a) + \delta(\omega + a))$                                      |
| 20 | $\sin(ax^2)$              | $-\sqrt{\frac{\pi}{a}} \sin\left(\frac{\omega^2}{4a} - \frac{\pi}{4}\right), a > 0$ |
| 21 | $\cos(ax^2)$              | $\sqrt{\frac{\pi}{a}} \cos\left(\frac{\omega^2}{4a} - \frac{\pi}{4}\right)$         |
| 22 | $\operatorname{sech}(ax)$ | $\frac{\pi}{a} \operatorname{sech}\left(\frac{\pi}{2a}\omega\right)$                |
| 23 | $e^{-\pi i a x^2}$        | $\frac{1}{\sqrt{a}} e^{-\frac{i\pi}{4}} e^{\frac{i\omega^2}{4\pi a}}, a > 0$        |
| 24 | $x^n$                     | $2\pi i^n \delta^{(n)}(\omega)$                                                     |
| 25 | $\delta^{(n)}(x)$         | $(i\omega)^n$                                                                       |
| 26 | $\frac{1}{x}$             | $-i\pi \operatorname{sgn}(\omega)$                                                  |
| 27 | $\frac{1}{x^n}$           | $-i\pi \frac{(-i\omega)^{n-1}}{(n-1)!} \operatorname{sgn}(\omega)$                  |
| 28 | $\frac{1}{\sqrt{ x }}$    | $\frac{\sqrt{2\pi}}{\sqrt{ \omega }}$                                               |

## Z-Transforms

| Technique                                                | Equations and Descriptions                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
|----------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Z-Transform</b>                                       | $\mathcal{Z}\{x[n]\} = X(z)$ <ul style="list-style-type: none"> <li>Converts a discrete sequence <math>x[n]</math> into a function <math>X(z)</math> of a complex variable <math>z</math>.</li> <li>Transforms linear difference equations with constant coefficients into algebraic equations.</li> <li>Used for recurrences, sampled signals, and digital systems.</li> <li>This sheet uses the one-sided (unilateral) Z-transform, with <math>n = 0, 1, 2, \dots</math></li> <li>The region of convergence (ROC) consists of the values of <math>z</math> for which the series converges absolutely.</li> </ul> |
| <b>Inverse Z-Transform</b>                               | $\mathcal{Z}^{-1}\{X(z)\} = x[n]$ <ul style="list-style-type: none"> <li>Use a transform table, partial fractions, or expansion in powers of <math>z^{-1}</math>.</li> <li>In the expansion, the coefficient of <math>z^{-n}</math> is <math>x[n]</math>.</li> </ul>                                                                                                                                                                                                                                                                                                                                               |
| <b>Z-Transform Equation</b>                              | $\mathcal{Z}\{x[n]\} = X(z) = \sum_{n=0}^{\infty} x[n] z^{-n}$ <p>where</p> <ul style="list-style-type: none"> <li><math>n</math>: integer sample index with <math>n = 0, 1, 2, \dots</math></li> <li><math>z</math>: dimensionless complex transform variable.</li> <li><math>u[n]</math>: unit step; equals 1 for <math>n \geq 0</math> and 0 otherwise.</li> <li><math>\delta[n]</math>: unit impulse; equals 1 at <math>n = 0</math> and 0 otherwise.</li> </ul>                                                                                                                                               |
| <b>Using Z-Transforms to Solve a Difference Equation</b> | <ol style="list-style-type: none"> <li>1. Apply the Z-transform to each term.</li> <li>2. Use the shift formulas and substitute the initial values.</li> <li>3. Solve the resulting algebraic equation for <math>Y(z)</math>.</li> <li>4. Apply the inverse Z-transform to find <math>y[n]</math>.</li> </ol>                                                                                                                                                                                                                                                                                                      |
| <b>Connection to Laplace Transforms</b>                  | <ul style="list-style-type: none"> <li>For sampling interval <math>T</math>, the substitution <math>z = e^{sT}</math> gives <math>z^{-n} = e^{-snT}</math>.</li> <li>This relates the transform kernels; a Z-transform is not generally equal to the Laplace transform of the original continuous-time signal.</li> </ul>                                                                                                                                                                                                                                                                                          |
| <b>Connection to Fourier Transforms</b>                  | <ul style="list-style-type: none"> <li>On the unit circle, <math>z = e^{i\Omega}</math>, the Z-transform equals the discrete-time Fourier transform when the unit circle lies in the ROC.</li> <li>Here, <math>\Omega</math> is the angular frequency in radians per sample.</li> </ul>                                                                                                                                                                                                                                                                                                                            |

**“Rabbit Growth”  
Example**

Let  $p[n]$  count rabbit pairs at year  $n$ , assuming no deaths and that each pair begins producing one new pair per year after a one-year maturation period.

Difference Equation:

$$p[n + 2] = p[n + 1] + p[n], \quad p[0] = 1, \quad p[1] = 2.$$

Define  $P(z) = Z\{p[n]\}$ . Applying the shift formulas:

$$z^2 P(z) - z^2 p[0] - z p[1] = z P(z) - z p[0] + P(z).$$

Therefore,

$$P(z) = \frac{z^2 + z}{z^2 - z - 1}.$$

Taking the inverse transform:

$$p[n] = \frac{1}{\sqrt{5}} \left[ \left( \frac{1 + \sqrt{5}}{2} \right)^{n+2} - \left( \frac{1 - \sqrt{5}}{2} \right)^{n+2} \right].$$

The sequence 1, 2, 3, 5, 8, 13, ... is the Fibonacci sequence starting at  $F_2$ .




**Table of One-Sided Z-Transforms**

| #  | Sequence<br>$x[n]$                      | Z-Transform<br>$X(z)$                                      | Region of<br>Convergence              |
|----|-----------------------------------------|------------------------------------------------------------|---------------------------------------|
| 1  | $\delta[n]$                             | 1                                                          | All $z$                               |
| 2  | $\delta[n - m]$<br>integer $m \geq 1$   | $z^{-m}$                                                   | $z \neq 0$                            |
| 3  | $u[n]$                                  | $\frac{z}{z - 1}$                                          | $ z  > 1$                             |
| 4  | $a^n$                                   | $\frac{z}{z - a}$                                          | $ z  >  a $                           |
| 5  | $n$                                     | $\frac{z}{(z - 1)^2}$                                      | $ z  > 1$                             |
| 6  | $n^2$                                   | $\frac{z(z + 1)}{(z - 1)^3}$                               | $ z  > 1$                             |
| 7  | $n^3$                                   | $\frac{z(z^2 + 4z + 1)}{(z - 1)^4}$                        | $ z  > 1$                             |
| 8  | $na^n$                                  | $\frac{az}{(z - a)^2}$                                     | $ z  >  a $                           |
| 9  | $n^2a^n$                                | $\frac{az(z + a)}{(z - a)^3}$                              | $ z  >  a $                           |
| 10 | $\cos(n\theta)$                         | $\frac{z(z - \cos \theta)}{z^2 - 2z \cos \theta + 1}$      | $ z  > 1$                             |
| 11 | $\sin(n\theta)$                         | $\frac{z \sin \theta}{z^2 - 2z \cos \theta + 1}$           | $ z  > 1$                             |
| 12 | $a^n \cos(n\theta)$                     | $\frac{z(z - a \cos \theta)}{z^2 - 2az \cos \theta + a^2}$ | $ z  >  a $                           |
| 13 | $a^n \sin(n\theta)$                     | $\frac{az \sin \theta}{z^2 - 2az \cos \theta + a^2}$       | $ z  >  a $                           |
| 14 | $u[n] - u[n - N]$<br>Integer $N \geq 2$ | $\sum_{k=0}^{N-1} z^{-k} = \frac{1 - z^{-N}}{1 - z^{-1}}$  | $z \neq 0$<br>value at $z = 1$ is $N$ |

**Conditions**

- All sequences are evaluated for  $n \geq 0$ .
- Let  $X(z) = \mathcal{Z}\{x[n]\}$ ,  $Y(z) = \mathcal{Z}\{y[n]\}$ ,  $a \neq 0$ , and  $\theta$  be real.
- If  $\theta$  is an integer multiple of  $\pi$ , the sine sequence is identically zero, so its transform is zero for all  $z$ .

## Properties and Shift Formulas for Z-Transforms

| #  | Property                               | Sequence / Operation      | Z-Transform                                      |
|----|----------------------------------------|---------------------------|--------------------------------------------------|
| 15 | Linearity                              | $Ax[n] + By[n]$           | $AX(z) + BY(z)$                                  |
| 16 | Delay by $m$ samples                   | $x[n - m] u[n - m]$       | $z^{-m}X(z)$                                     |
| 17 | Advance by one sample                  | $x[n + 1]$                | $zX(z) - zx[0]$                                  |
| 18 | Advance by two samples                 | $x[n + 2]$                | $z^2X(z) - z^2x[0] - zx[1]$                      |
| 19 | Advance by $m$ samples                 | $x[n + m]$                | $z^mX(z) - \sum_{k=0}^{m-1} x[k]z^{m-k}$         |
| 20 | Exponential multiplication             | $a^n x[n]$                | $X(z/a)$                                         |
| 21 | Multiplication by $n$                  | $nx[n]$                   | $-z \frac{dX}{dz}$                               |
| 22 | Running sum                            | $\sum_{k=0}^n x[k]$       | $\frac{z}{z-1}X(z)$                              |
| 23 | Convolution                            | $\sum_{k=0}^n x[k]y[n-k]$ | $X(z)Y(z)$                                       |
| 24 | Periodic sequence<br>period $N \geq 1$ | $x[n + N] = x[n]$         | $\frac{\sum_{k=0}^{N-1} x[k]z^{-k}}{1 - z^{-N}}$ |

### Conditions

- In shift formulas,  $m$  is a positive integer.
- Delays use zero extension for negative indices.
- Linearity and convolution hold in the common ROC; cancellations may enlarge it.
- Running sums converge at least where  $X(z)$  converges and  $|z| > 1$ .
- For exponential multiplication, require  $z/a$  in the ROC of  $X$ .
- Differentiation holds inside the ROC.
- A nonzero periodic sequence has ROC  $|z| > 1$ .

### Initial- and Final-Value Theorems

| Theorem       | Equation                                                                     | Conditions                                                                                                       |
|---------------|------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------|
| Initial Value | $x[0] = \lim_{z \rightarrow \infty} X(z)$                                    | The one-sided transform converges for sufficiently large $ z $ .                                                 |
| Final Value   | $\lim_{n \rightarrow \infty} x[n] = \lim_{z \rightarrow 1} (1 - z^{-1})X(z)$ | For rational $X(z)$ , all poles of $(1 - z^{-1})X(z)$ , after cancellation, must lie strictly inside $ z  < 1$ . |

## Sources

- Dawkins, Paul (2004). Table of Laplace Transforms. [https://tutorial.math.lamar.edu/classes/de/Laplace\\_Table.aspx](https://tutorial.math.lamar.edu/classes/de/Laplace_Table.aspx)
- Furius Enterprises (2010). Differential Equations Cheat Sheet 2nd-order Homogeneous. <https://furius.ca/cgfpub/doc/diffequs/diffequs.pdf>
- Shapiro, B.E. (2014). Table of Laplace Transforms. <https://www.integral-table.com/downloads/LaplaceTable.pdf>
- MathWorks: Solve Difference Equations Using Z-Transform. [https://www.mathworks.com/help/symbolic/compute-z-transforms-and-inverse-z-transforms.html?utm\\_source=chatgpt.com](https://www.mathworks.com/help/symbolic/compute-z-transforms-and-inverse-z-transforms.html?utm_source=chatgpt.com)
- Wikipedia (2025). Tables of important Fourier transforms. [https://en.wikipedia.org/wiki/Fourier\\_transform#Tables\\_of\\_important\\_Fourier\\_transforms](https://en.wikipedia.org/wiki/Fourier_transform#Tables_of_important_Fourier_transforms)

## See Also

- [Harold's Partial Differential Equations \(PDE\) Cheat Sheet](#)
- [Harold's Differential Equation Models Cheat Sheet](#)
- [Harold's Euler's Method Example](#)

## Acknowledgements

- Proofread with assistance from ChatGPT 6.1 Sol, <https://chatgpt.com/>